

# Optimizing Curbside Parking Resources Subject to Congestion Constraints

Chase Dowling, Tanner Fiez, Lillian Ratliff, and Baosen Zhang<sup>‡</sup>

March 22, 2017

## Abstract

To gain theoretical insight into the relationship between parking scarcity and congestion, we describe block-faces of curbside parking as a network of queues. Due to the nature of this network, canonical queueing network results are not available to us. We present a new kind of queueing network subject to customer rejection due to the lack of available servers. We provide conditions for such networks to be stable, a computationally tractable “single node” view of such a network, and show that maximizing the occupancy through price control of such queues, and subject to constraints on the allowable congestion between queues searching for an available server, is a convex optimization problem. We demonstrate an application of this method in the Mission District of San Francisco; our results suggest congestion due to drivers searching for parking stems from an inefficient spatial utilization of parking resources.

## 1 INTRODUCTION

Drivers in densely populated urban districts often find that desirable parking close to their destination is unavailable or prohibitively expensive. Drivers will begin to *cruise for parking* [Shoup, 2005], significantly contributing to surface street congestion. Researchers have attempted to measure the economic loss to both these drivers and the cities themselves. For the former, drivers in different cities can spend anywhere between 3.5 to 14 minutes searching for spots every time they park [Shoup, 2006]. For the latter, cruising behaviors can lead to substantial congestion in dense urban districts. For instance, there exists a commonly cited folklore that 30% of traffic in a city is directly due to drivers looking for parking [Shoup, 2005]<sup>1</sup>.

Municipalities and city planners typically aim to achieve some target *occupancy*: the percentage of parking spaces in use at any given time [SFpark, 2013]. Fig. 1 shows the

---

\*The authors have been supported in part by NSF grants CNS-1646912 and CNS-1634136. C. Dowling was also supported in part by the Washington Clean Energy Institute.

<sup>†</sup>C. Dowling, T. Fiez, L. Ratliff and B. Zhang are with the Department of Electrical Engineering, University of Washington, Seattle, WA 98195, USA Emails: {cdowling,fiezt,ratliff,zhangbao}@uw.edu

<sup>1</sup>It is not entirely clear where this number originates from as estimates vary wildly. A motivation of this paper is to rigorously quantify the actual effect of parking on congestion.

occupancy of the 3400 block of 18th St. in San Francisco, CA. Cities like San Francisco have launched projects like SFPark to target an average occupancy between around 85% by slowly adjusting prices based on observed demand [SFPark, 2013].




Fig. 1: The observed parking occupancy (percentage of parking spaces in use), on a Saturday along the 3400 block of 18th St. in the Mission District of San Francisco. Fig. 2 illustrates a key result of this paper: the congestion resulting from lack of parking along a block-face.

Parking *occupancy* (and availability) is an indirect measure (and means of control) of overall demand for vehicle access. Yet, if city planners must control congestion, occupancy alone is not a sufficient measure. Firstly, the same occupancy levels of two streets in different parts of the city can lead to different effects on through-traffic delays or respond differently to incremental price changes. Secondly, the street topology and interactions between different blocks can lead to complex traffic dynamics, which a single number like occupancy cannot capture. At the same time, cities cannot be overly aggressive in controlling parking occupancy since they must maintain a high availability of parking resources to serve downtown businesses and residents, as well as delivery, courier, and emergency vehicle services. Therefore, a reasonable question that a city planner would be interested in addressing is the following: *Given a maximum tolerable level of congestion, what is the maximum occupancy at a block and what price achieves this occupancy?*

The question of parking’s impact on congestion has remained difficult to address due to: 1) lack relevant data on pricing and demand and 2) lack of tractable and rigorous models that link parking to congestion and capture spatial and temporal variation. To address this question utilizing parking occupancy, traffic, and surface street topology data that is available today, our contributions are:

1. *Modeling*: we describe and analyze a new kind of queue network where customers move between queues according to a network topology *until* an available server is found, and leave the network after service
2. *Control*: we show that maximizing occupancy subject to constraints on the congestion created by drivers searching for parking is a convex program.
3. *Application*: we conduct a study based on real occupancy and pricing data for blocks in the San Francisco Mission District, showing that a) higher total occupancy does not necessarily lead to more traffic, and b) incentivizing drivers to park further away by




Fig. 2: A visualization of one of our key results: estimated percentage of through traffic searching for parking. This estimate is obtained by determining the minimum arrival rate necessary to achieve an observed occupancy (Fig. 1), and comparing this to the observed total through-traffic.

reducing price can be equally as effective as disincentivizing drivers from parking at desirable locations.

The paper is organized as follows. We provide motivation and review related work in Section 2. In Section 3, we present the network queue model. We present results in Section 4. In particular, we provide stability conditions under a uniformity assumption on the network topology, we provide a framework for determining the arrival rate in the non-uniform case, and we pose an optimization problem to optimize parking availability subject to maximum congestion constraints that we show to be convex. In Section 6, we demonstrate the effectiveness of the solution to the optimization problem on a network modeled after San Francisco’s Mission District. We conclude with discussion and commentary on future work in Section 7.

## 2 Motivation

As observed by Pierce and Shoup, circling for parking occurs when occupancy reaches 100% [Pierce and Shoup, 2013], however, this takes an instantaneous point of view likely unavailable to city planners. Rather, if occupancy is taken to be the expected proportion of parking spaces in use *over a given time period*, then high occupancy block-faces must be full at least some of the time, and therefore responsible for some traffic—see, *e.g.*, Fig. 2.

### 2.1 Data Availability

Municipalities (in particular, city planners and transportation departments) are gaining access to data from recently installed *smart parking meters* and, on occasion, individual parking space sensors (e.g., San Francisco [SFpark, 2013], Seattle [Ottosson et al., 2013], Los Angeles [Ghent et al., 2012], and Pittsburgh [Fabusuyi et al., 2013]). Yet, no city has completely implemented full-scale transportation sensor grids that include active monitoring of parking on a space-by-space basis. Regardless whether such a goal may be reached, however, many cities have a growing history of parking transaction data collected by digital

meters. These data can be used to estimate parking occupancy; transactions provide an estimate of how long a driver intended to park and the number of drivers parked moment to moment. In our experiments, we make use of transaction, traffic, and infrastructural data publicly made available by the SFPark pilot study [Agency, 2017].

## 2.2 Related Work

Early work focused largely on parking supply and demand [Vickrey, 1954], and refinement of the economic view of parking continues through today [Inci, 2015]. The costs of congestion caused by *cruising for parking* [Shoup, 2005, Shoup, 2006] have motivated research in modeling urban parking dynamics, and economizing of parking spaces has led to a desire to control demand levels via price.

Over the last few decades, a number of models (e.g., Vickrey’s celebrated “bathtub” model) have been developed and introduced in the absence of data only recently becoming available [Arnott, 2013, Arnott and Inci, 2006]. These models typically take a time-varying flow and capacity view in the form of systems of partial differential equations (see [Inci, 2015] for an overview of variations on these models).

Recent research has observed, however, that transaction data can be used to estimate parking occupancy and, in consequence, used to estimate resource performance [Yang and Qian, 2017, Dowling et al., 2017]. The distinction that occupancy below 100% results in congestion has recently been noted by [Millard-Ball et al., 2014] in their own analysis of the SFPark pilot study parallel to [Pierce and Shoup, 2013], however the authors of [Millard-Ball et al., 2014] view block-face parking as a Bernoulli random variable, between being full or not. We build on this work by 1) *not* implicitly assuming curbside parking occupancy is independent between block-faces and 2) considering all possible states of parking spaces—from completely empty to completely full—along block-faces.


Occupancy and other data lend themselves to discrete and probabilistic models that may potentially better reflect flow on surface streets as compared to flow on highways or through spatially homogeneous regions, as in [Arnott, 2006] and [Arnott and Rowse, 2009]. Hence, classical methods of queueing theory have recently been applied to parking areas: garage and curbside alike [Ratliff et al., 2016, Larson and Sasanuma, 2010, Caliskan et al., 2007, Ceballos and Curtis, 2004].

Our work primarily builds on existing parking literature by expressing curbside parking as a network of queues. Specifically, utilizing newly available parking data, we implement the basis for a spatially heterogeneous model city planners can use to effectively test parking policies and, furthermore, we determine that maximizing occupancy subject to congestion constraints using price controls is a convex optimization problem.

## 3 Queueing Model

### 3.1 Model Setup

Although a natural model, queue networks have not been used extensively in parking related research (see, e.g. [Raheja, 2010] and the references within for more details). Two major



(a) Example of block-face adjacency with respect to one-way (green, one arrow) and two-way (red, two arrows) streets. The corresponding network version is illustrated by Fig. 3b.

(b) Corresponding graphical representation of Fig. 3a with respect to individual sides of a street. The solid arrows are edges between block-faces visible in Fig. 3a, while the dashed arrows are between block-faces not labeled. Note that drivers leaving the red, two-way block-faces (1 and 3) may only continue straight or turn right, while drivers leaving the green, one-way block-faces (2 and 4) can continue straight, or turn right or left.

Fig. 3: Block-face parking around a typical city block, and the corresponding graphical representation of the queue-network with respect to legal turns.

reasons for this are: 1) the size of the state space grows exponentially as the size of the network grows; 2) established queueing network results (e.g., for communication networks) do not carry over directly. The rest of this section will describe the details of the queueing network model, its difference to conventional models, and how we overcome these difficulties.

### 3.2 Queues Interacting Via Rejections

We model each block-face as a multi-server queue, where the number of servers is the number of available parking spots on that block-face. The block-faces are connected as *nodes on a graph*, where two nodes are adjacent if vehicles can go from one block-face to the other in the road network. See Fig. 3a for an example. To account for legal turning maneuvers (e.g., right turn only) and one way streets, we use *directed edges*. We use conventional notations  $D = (V, E)$  to describe this digraph. Without loss of generality (WLOG), we assume this graph is connected.

A queue, or a node  $i \in V$  is characterized by an *exogenous* arrival rate  $\lambda_i$ , a service rate  $\mu_i$ , and the number of servers  $k_i$ . We assume that the *exogenous* arrival process is Poisson (independent between queues) and the services times are generally distributed like conventional  $M/G/\cdot/\cdot$  queues [Wolff, 1989], however, unlike conventional queueing networks where customers are buffered at individual queues, we assume that customers (or drivers), are buffered or queued along the *network edges*. This behavior reflects the key fact that vehicles which cannot find parking circulate in the network rather than wait at one location. Therefore, if the driver is served by a queue, it then leaves the network. However, if it finds

the current queue to be full, it is *rejected by that queue* and moves to neighboring queues to find new parking spots. The rate of these rejections is parking scarcity’s contribution to through-traffic delays.

The key difference between our queue network and conventional networks—such as a Jackson network [Jackson, 1957]—is that drivers proceed to other queues after they are rejected rather than served. Since the rejection of a queue with Poisson arrivals and exponential service times is not Poisson, characterizing the stationary distribution of this network of queues is very difficult because the distribution of total arrival rate itself to any queue is unknown.

Since the exact distribution of the queue is difficult to characterize, we instead turn to understanding the behavior of the *mean performance metrics* of the network. This relaxation allows us to use theorems such as Little’s Law [Little, 1961] that do not depend on the exact distributions. Secondly, the controllable and measurable quantities are often average values like occupancy and parking service times.

### 3.3 Stationary Distribution of a Single Queue

Here we introduce how a *single queue* can be analyzed, and later in the paper extend the analysis to a network of queues. To help avoid confusion between exogenous arrivals (from outside of the network, denoted by  $\lambda$ ) and endogenous arrivals (rejection from neighboring queues, denoted by  $x$ ), we use  $y$  as the total arrival rate to a queue. Suppose the service rate (inverse length of parking time) of each server is  $\frac{1}{\mu}$  and there are  $k$  servers ( $k$  parking spots) in total. Let  $\pi_i$  be the stationary probability that  $i$  servers are busy ( $i$  cars are parked), for  $i = 0, \dots, k$ . Let  $\boldsymbol{\pi} = [\pi_0 \dots \pi_k]$ . For this single queue, we can explicitly write down its stationary probability distribution via the transition rate matrix:

$$\mathbf{Q} = \begin{bmatrix} -y & y & 0 & 0 & \dots & 0 & 0 \\ \mu & -(\mu + y) & y & 0 & \dots & 0 & 0 \\ 0 & 2\mu & -(2\mu + y) & y & \dots & 0 & 0 \\ & & & & \vdots & & \\ 0 & 0 & 0 & 0 & \dots & k\mu & -k\mu \end{bmatrix},$$

and  $\boldsymbol{\pi}$  is the unique solution to

$$\boldsymbol{\pi}\mathbf{Q} = \mathbf{0} \tag{1}$$

such that  $\sum \pi_i = 1$ . Let  $\rho = \frac{y}{\mu}$ . By standard calculations [Wolff, 1989],

$$\boldsymbol{\pi} = \pi_0 \cdot \left[ 1, \rho, \dots, \frac{\rho^k}{k!} \right] \tag{2}$$

where  $\pi_0 = \left[ \sum_{j=0}^k \frac{\rho^j}{j!} \right]^{-1}$ . Using Little’s Law, the occupancy  $u$ , or the proportion of busy servers at any given time can be expressed as,

$$u = \frac{y}{k\mu} \left( 1 - \pi_0 \frac{\rho^k}{k!} \right) \tag{3}$$

Note that  $(1 - \pi_0 \frac{y^k}{k!})$  is the probability that *at least* one space is available. Consider, if drivers are unable to wait for an available server at a particular block, in order to obtain occupancies approaching 100%, cars would need to arrive at an infinite rate in order to immediately replace vehicles exiting service. Since it is often cited that congestion due to driver's searching for parking is a significant cost to the social welfare, this is a critical misconception.

A block-face queue is therefore rejecting incoming vehicles at a rate of  $y \cdot \pi_k$ . The difficulty therein lies with estimating these total arrival rates, because no two adjacent block-faces are independent.

## 4 Network of Queues

In this section we study these networks of queues. We first consider the uniform case, then extend the results to the non-uniform case.

### 4.1 Uniform Network

Many urban centers have fairly uniform street topologies (e.g., the famed Manhattan streets), where the streets form a regular graph. In this section we make the assumption that the queueing network is entirely uniform: the topology is a  $d$ -regular graph, all block-faces have the same number of servers with the same service rate  $\mu$ , and they have the same exogenous arrival rate  $\lambda$ .

In this regular queue network, each queue will have equal stationary distributions in the steady state, therefore we only need to look at a single queue as representative of the state space of the entire network. Let  $x$  be the average rate of rejection of a queue to one of its neighbors, and  $dx$  be the total rejection to all of its neighbors. Let  $y = \lambda + dx$  be the total arrival rate to a queue, where  $\lambda$  is the exogenous arrivals and  $dx$  are the rejections from its neighboring queues. We have the conservation equation,

$$dx = y\pi_k, \tag{4}$$

where  $\pi_k$  is the probability that all  $k$  servers are busy. Combined with stationary distribution of (1) we have the following equations:

$$\begin{cases} \pi Q & = 0 \\ \sum \pi_i & = 1 \\ dx & = \pi_k(\lambda + dx) \end{cases} \tag{5}$$

We can write (4) as,

$$y - \lambda = \frac{\rho^k}{k!} y \tag{6}$$

where  $\rho = \frac{y}{\mu}$ . The equation in (6) is a polynomial in  $y$ . The next lemma states that there exists a unique solution to  $y$  (and thus  $x$ ) as long as the queues are stable:

**Lemma 1.** *If  $0 < \lambda < \mu k$ , then (i) there is a unique and positive solution to  $y$  in (6) and (ii) the solution is greater than  $\lambda$ . In addition, the rejection rate  $x$  is also unique and positive.*

The proof is given in Appendix .1. This result states that as long as the total arrivals are less than the service rate times the number of spaces, we can explicitly find the rejection rates and the stationary probabilities by solving a polynomial equation.

## 4.2 Non-uniform Network

Of course, the totally uniform assumption rarely holds up in practice. But given occupancy data we show that the *total* exogenous and endogenous arrivals to a queue can still be solved for and used to estimate the traffic caused by drivers searching for parking. This time, for some *total* incoming rejection rate  $x$ , letting  $y = \lambda + x$ , we can estimate the endogenous proportion of incoming arrivals as the sum of the outgoing fractional rejection rates of adjacent queues.

Assuming the queueing network reaches steady state, from the perspective of a single queue in solving 3 for  $\pi_0$  gives

$$\pi_0 \frac{\rho^k}{k!} + \frac{uk\mu}{y} = 1, \quad (7)$$

where  $u$  is the occupancy level and  $\rho = \frac{y}{\mu}$ . Rearranging terms yields a polynomial in  $y$ ,

$$0 = \sum_{i=0}^k \frac{1}{\mu^{i-1}} \left[ \frac{i - uk}{i!} \right] y^k. \quad (8)$$

Again, we can characterize the solutions to (8)

**Lemma 2.** *If  $u \in [0, 1)$  and  $k$  is a positive integer, then (8) has a unique real, positive root.*

The proof is provided in Appendix .2.

This root need not be bounded, hence the restriction of the values of  $u$  to the interval  $[0, 1)$ . In order to achieve a 100% occupancy, implying the probability of being full is 1, vehicles would need to arrive constantly ( $y = \infty$ ), immediately taking the place of any vehicle that leaves upon completion of service. This is analogous to the requirement that for the  $M/M/k/k$  queue to be stable,  $\pi_0 > 0$ .

## 5 Optimizing parking availability

Price elasticity of demand provides a means of describing how consumer demand will change with incremental changes to price. Currently, Pierce and Shoup's analysis of the SFPark pilot project in [Pierce and Shoup, 2013] is the state-of-the-art in estimating the price elasticity of demand for curbside parking; their exploratory analysis provided rough estimates of aggregated elasticities across time, location, and price change directions. For the purposes of this paper, and in order to make use of the results in [Pierce and Shoup, 2013] we assume a *linear* elasticity, however, any demonstrably reasonable (reflective of consumer behavior),



concave function would not tax the validity of our results. Thus, a block-face has a linear elasticity  $\alpha$ , and a function  $\mathcal{U} : p \mapsto u$ , taking a price  $p$  to an occupancy level  $u$ , defined as

$$\mathcal{U}(p) = 1 - \alpha p \tag{9}$$

Recall (8); we can write the right-hand side of this equation as a mapping  $F : Y \times U \rightarrow \mathbb{R}$  where  $U = (0, 1)$  such that

$$F(y, u) = \sum_{i=0}^k \frac{1}{\mu^{i-1}} \left[ \frac{i - uk}{i!} \right] y^k \tag{10}$$

Note that this map is smooth in both its arguments  $y$  and  $u$ . By applying the Implicit Function Theorem [Lee, 2012, Theorem C.40], a smooth mapping  $f : u \mapsto y$  exists and it is continuous and differentiable. Moreover, there is an explicit expression for its derivative and the function  $f$  maps an occupancy  $u \in U$  to the unique real root  $y$  of  $F(y, u) = 0$ .

Consider the following composition,

$$g(p) = f(\mathcal{U}(p)) \cdot \pi_k, \tag{11}$$

which is equal the rate of rejection of vehicles from a block given a price  $p$ . The composition (11) takes a price to a resulting level of congestion along an edge in a queue network due to rejections.

The optimization problem given by

$$\begin{aligned} & \underset{\mathbf{p}}{\text{maximize}} && \mathcal{U}(\mathbf{p}) \\ & \text{subject to} && g_i(p_i) \leq \bar{x}_i, \quad i = 1, \dots, m. \end{aligned} \tag{P-1}$$

maximizes parking resource utilization subject to a congestion constraints  $\bar{x}_i$  imposed on each block-face. Since (9) is concave, if  $g_i$ 's are convex, then (P-1) is a convex optimization problem easily solved by gradient descent.

**Theorem 1.** *The optimization problem (P-1) is convex.*

The main technical challenge is to show that the constraints  $g_i$ 's are convex in the occupancy  $u_i$ . This is somewhat involved and the proof is provided in Appendix .3. Since (P-1) is convex, there exist many ways to solve it. We use a projected gradient for the case study in the next section.

## 6 APPLICATION

We consider the application of the above methods to curbside parking San Francisco's Mission District (Fig. 4). Using data collected by the SFPark pilot from May 8th, 2012 - August 29th, 2017 and elasticities estimated by [Pierce and Shoup, 2013], we identify block-faces responsible for the high congestion impacts to through-traffic and set constraints to bring this down to some hypothetically tolerable level. All data is calculated to an *hourly* rate, e.g. the average percentage of parking spaces in use over the course of an hour.

According to [Pierce and Shoup, 2013], curbside parking in the Mission District of San Francisco displayed an average price elasticity of  $-0.21$ . Price elasticity varied greatly due to the time of day, week, and year, among a number of other observable factors. For the purposes of demonstration in this paper, we assume a uniform price elasticity of  $-0.21$  across block-faces in the Mission District, and therefore, resulting price changes should be taken with a grain of salt.

We examine two scenarios: 1) we wish to reduce overall *congestion* due to parking by 80% at two high occupancy block-faces and 2) achieve  $>80\%$  *occupancy* at each block-face, rather than a neighborhoodwide average of 80%, concentrated at a smaller proportion of the blocks in the district. We find that, in particular, *spatial inefficiency, and not high occupancy, results in congestion*.

### 6.0.1 Congestion Reduction

The 3300 block of 17<sup>th</sup> street and the 3400 block of 18<sup>th</sup> street are responsible for the overwhelming majority of parking related congestion in Mission District at noon on the average Saturday, generating a total of nearly 60 vehicles unable to find parking per hour. As illustrated by Fig. 2, a full third of 18<sup>th</sup> street’s through traffic is made up of drivers unable to find parking.

At these traffic levels, 17<sup>th</sup> and 18<sup>th</sup> street have occupancies of 97% and 98% respectively. By increasing prices by \$0.28 on 17<sup>th</sup> and \$0.27 on 18<sup>th</sup>, we are able to reduce this congestion by 80% to approximately 11 vehicles per hour, total, while still maintaining 91% and 92% occupancies, respectively. All other blocks see comparatively negligible changes.

The “elbow” of the highly non-linear curve describing the total arrival rate needed to achieve a particular occupancy level occurs around the 90% mark, as illustrated in




Fig. 4: Block-faces, highlighted in red, with curbside parking data from the SFPark pilot program; Mission District, San Francisco.





Fig. 5: Necessary total arrival rate  $y$  to achieve an occupancy level for some fixed number of servers  $k$  with a service time  $\mu$  of 1. Not the sharp increase in total arrival rate around the 90% occupancy mark and that increasing the number of servers only has a marginal bearing on this arrival rate.

Fig. 5. By redistributing vehicles intending to park at high occupancy blocks to historically low occupancy blocks through price control, less time is spent cruising for parking, leading us to our next experiment.


### 6.0.2 Occupancy Redistribution

On a typical Saturday at noon, the Mission District achieves an average occupancy of approximately 78%, while generating over 60 vehicles per hour in additional traffic due to drivers searching for parking because there is a small number of high occupancy block-faces and a larger number of low occupancy block-faces. By bounding each block to producing no more than 1 vehicle every 20 minutes unable to find parking (for a total of 48 per hour for the district), each individual block-face individually exceeds 85% occupancy *at each block-face*. Indeed, after price control, the Mission District services a larger *total* number of vehicles while still producing less additional traffic due parking scarcity.


Fig. 6c indicates that, significantly discounting prices on low occupancy block-faces is an equally effective solution as raising prices at high occupancy block-faces, in order to achieve an effective distribution of parking resources that does not generate a costly amount of congestion searching for parking. Indeed, considering that a small number of block-faces may exhibit a high occupancy due to their desirable proximity to popular locations, incentivizing drivers to park somewhat further away may be more effective than pricing out other drivers by means of money or time to walk to a location.



(a) Occupancy and resulting traffic in vehicles per hour generated.



(b) Redistributing demand in Fig. 6a to low-occupancy block-faces using the price changes indicated in Fig. 6c results in less total traffic.



(c) Price changes corresponding to the resulting occupancy redistribution in 6.0.2

Fig. 6: Results of experiments in section 6.0.2

## 7 CONCLUSIONS AND FUTURE WORKS

### 7.1 Conclusions

With the growth of ride sharing services, electric vehicles, and increased demand for local delivery services, personal and commercial transportation is changing. In order for city plan-

ners to design effective future parking policies and make use of growing bodies of parking data, we developed a new kind of queueing network. We provided conditions for such networks to be stable, a “single node” view of a queue in such a network, and showed that maximizing the occupancy of such queues subject to constraints on the allowable congestion between queues searching for an available server is a convex optimization problem.

## 7.2 Future Works

A standing question in parking economics research is that of an appropriate maximum parking time [Inci, 2015]. Some argue that a lower maximum parking time or lack of an initial buy-in price results in higher vehicle turn-over, and hence more congestion. Indeed, according to (3), decreasing  $\mu$  increases the total arrival rate necessary to achieve a fixed occupancy, but the probability of being full remains unchanged. Combined with the collection of ground-truth data and hypothesis testing, this question is closer to being answered.

Further, driver behavior is an important next-step to be considered. We have implicitly assumed that drivers, once inside the network searching for parking, will park regardless of price at a particular block-face. While this assumption alone is not unrealistic, how demand changes with respect to the total network sojourn time of the driver, distance from the initially desired location, and whether or not drivers have access to information regarding available parking locations are all certainly critical implications to consider.

## References

- [Agency, 2017] Agency, S. F. M. T. (2017). Sfpark. <http://sfpark.org/>.
- [Arnott, 2006] Arnott, R. (2006). Spatial competition between parking garages and downtown parking policy. *Transport Policy*, 13(6):458–469.
- [Arnott, 2013] Arnott, R. (2013). A bathtub model of downtown traffic congestion. *Journal of Urban Economics*, 76:110–121.
- [Arnott and Inci, 2006] Arnott, R. and Inci, E. (2006). An integrated model of downtown parking and traffic congestion. *J. Urban Economics*, 60(3):418–442.
- [Arnott and Rowse, 2009] Arnott, R. and Rowse, J. (2009). Downtown parking in auto city. *Regional Science and Urban Economics*, 39(1):1–14.
- [Caliskan et al., 2007] Caliskan, M., Barthels, A., Scheuermann, B., and Mauve, M. (2007). Predicting parking lot occupancy in vehicular ad hoc networks. In *Vehicular Technology Conference, 2007. VTC2007-Spring. IEEE 65th*, pages 277–281. IEEE.
- [Ceballos and Curtis, 2004] Ceballos, G. and Curtis, O. (2004). Queue analysis at toll and parking exit plazas: a comparison between multi-server queueing models and traffic simulation. In *ITE Annual Meeting and Exhibit*.
- [Dowling et al., 2017] Dowling, C., Fiez, T., Ratliff, L., and Zhang, B. (2017). How much urban traffic is searching for parking? *arXiv preprint arXiv:1702.06156*.

- [Fabusuyi et al., 2013] Fabusuyi, T., Hampshire, R., and Hill, V. (2013). Evaluation of a smart parking system. *Transportation Research Record: Journal of the Transportation Research Board*, (2359):10–16.
- [Ghent et al., 2012] Ghent, P., Mitchell, D., and Sedadi, A. (2012). La express park<sup>TM</sup>-curbing downtown congestion through intelligent parking management. In *19th ITS World Congress*.
- [Inci, 2015] Inci, E. (2015). A review of the economics of parking. *Economics of Transportation*, 4(1):50–63.
- [Jackson, 1957] Jackson, J. (1957). Networks of waiting lines. *Operations Research*, 5(4):518–521.
- [Larson and Sasanuma, 2010] Larson, R. and Sasanuma, K. (2010). Congestion pricing: A parking queue model. *J. Industrial and Systems Engineering*, 4(1):1–17.
- [Lee, 2012] Lee, J. M. (2012). *Introduction to Smooth Manifolds*. Springer.
- [Little, 1961] Little, J. D. (1961). A proof for the queuing formula:  $L = \lambda w$ . *Operations research*, 9(3):383–387.
- [Meserve, 1982a] Meserve, B. E. (1982a). *Fundamental Concepts of Algebra*. Dover Publications.
- [Meserve, 1982b] Meserve, B. E. (1982b). *Fundamental Concepts of Algebra*. Dover Publications.
- [Millard-Ball et al., 2014] Millard-Ball, A., Weinberger, R. R., and Hampshire, R. C. (2014). Is the curb 80% full or 20% empty? assessing the impacts of san francisco’s parking pricing experiment. *Transportation Research Part A: Policy and Practice*, 63:76–92.
- [Ottooson et al., 2013] Ottooson, D. B., Chen, C., Wang, T., and Lin, H. (2013). The sensitivity of on-street parking demand in response to price changes: A case study in seattle, wa. *Transport Policy*, 25:222–232.
- [Pierce and Shoup, 2013] Pierce, G. and Shoup, D. (2013). Getting the prices right: an evaluation of pricing parking by demand in san francisco. *Journal of the American Planning Association*, 79(1):67–81.
- [Raheja, 2010] Raheja, T. (2010). Modelling traffic congestion using queuing networks. *Sadhana*, 35(4):427–431.
- [Ratliff et al., 2016] Ratliff, L., Dowling, C., Mazumdar, E., and Zhang, B. (2016). To observe or not to observe: Queuing game framework for urban parking. In *Proc. 55th IEEE Conference on Decision and Control*, pages 5286–5291.
- [SFpark, 2013] SFpark (2013). San francisco parking pilot evaluation.

[Shoup, 2005] Shoup, D. (2005). *The high cost of free parking*, volume 7. Planners Press, American Planning Association.

[Shoup, 2006] Shoup, D. (2006). Cruising for parking. *Transport Policy*, 13(6):479–486.

[Vickrey, 1954] Vickrey, W. (1954). The economizing of curb parking space. *Traffic Engineering*, 29(1):62–67.

[Wolff, 1989] Wolff, R. W. (1989). *Stochastic modeling and the theory of queues*. Pearson College Division.

[Yang and Qian, 2017] Yang, S. and Qian, Z. S. (2017). Turning meter transactions data into occupancy and payment behavioral information for on-street parking. *Transportation Research Part C: Emerging Technologies*, 78:165–182.

## .1 Proof of Lemma 1

*Proof.* Some algebra on (6) gives

$$k!(y - \lambda) \sum_{i=0}^k \frac{\rho^i}{i!} = y\rho^k$$

The  $\frac{y^{k+1}}{\mu^k}$  and  $y\rho^k$  terms cancel, and we have a polynomial with degree  $k$

$$\frac{\frac{k}{\mu^{k-1}} - \frac{\lambda}{\mu^k}}{k!} y^k + \frac{\frac{k-1}{\mu^{k-2}} - \frac{\lambda}{\mu^{k-1}}}{(k-1)!} y^{k-1} + \dots + \left(1 - \frac{\lambda}{\mu}\right) y - \lambda = 0. \quad (12)$$

Descartes' rule of signs [Meserve, 1982a], which roughly states that given a polynomial and ordering its terms from highest degree to lowest degree, the number of real positive roots is related to the number of sign changes. Let  $n$  be the number of sign changes (from positive to negative), then the only possible number of positive roots to this polynomial are  $n, n-2, n-4, \dots$ . In particular, if  $n = 1$ , then the polynomial has one and only one positive root. Applying to the polynomial in (12), we notice the sign of the coefficients are determined by  $k - \lambda$ ,  $k - 1 - \lambda$ ,  $k - 2 - \lambda$  and so on, until the constant term  $-\lambda$ . By assumption,  $\lambda < \mu k$ , so the first coefficient is positive. By assumption,  $\lambda > 0$ , so the last coefficient (constant term) is negative. Then for any  $\lambda \in (0, \mu k)$ , it causes at most one change the signs of the other coefficients. So  $n = 1$  for all possible  $\lambda \in (0, k)$ , and there is a unique positive solution to  $y$ .

To show that  $y > \lambda$ , let  $f(y)$  be the polynomial in (12). We have  $f(0) = -\lambda < 0$ , and  $f(z) > 0$  for sufficiently large  $z$  (positive coefficient on  $y^k$  term). Since there is only one positive solution, it suffices to show that at  $f(\lambda) < 0$ . It turns out that  $f(\lambda)$  has a telescoping sum, and

$$\begin{aligned} f(\lambda) &= \sum_{i=1}^k \frac{\lambda^i}{(i-1)!} - \sum_{i=1}^k \frac{\lambda^{i+1}}{i!} - \lambda \\ &= \lambda - \frac{\lambda^{k+1}}{k!} - \lambda \\ &< 0. \end{aligned}$$

□

## .2 Proof of Lemma 2

*Proof.* Let us first examine the coefficients of  $y^k$ . WLOG, assume  $\mu = 1$ . We have the following sequence:

$$s = \{-uk, 1 - uk, \frac{2-uk}{2!}, \dots, \frac{k-uk}{k!}\} \quad (13)$$

We will show that if  $u \in [0, 1)$ ,  $k \in \mathbb{Z}_+$ , the sequence (13) undergoes exactly 1 sign change, and again apply Descartes' rule of signs. Observe that  $s_0 < 0$  for any allowable values of  $u$  and  $k$ . Further, observe that  $s_k = (1 - u)((k - 1)!)^{-1}$ . By induction,  $s_k$  will always be positive for any value of  $k$ . If  $k = 1$ , then  $s_1 = (1 - u)(1)^{-1}$ , and since  $u \in [0, 1)$ ,  $s_1 > 0$ . Assume this is true for  $k$ , then for  $k + 1$ ,  $s_k = (1 - u)(k!)^{-1}$ , so that we have that  $s_{k+1} > 0$ . It now suffices to show that  $\{s\}$  can only undergo one sign change as we increment  $i$ . For some  $k$ , the  $i$ -th element of  $\{s\}$  is  $s_i = (i - uk)(i!)^{-1}$ . Fix  $k$ . While the denominator of the sequence is itself increasing with  $i$  (meaning  $\{s\}$  need not be monotonic), it is strictly positive. We need only look at the sign of the numerator. In particular,  $uk$  is fixed between  $[0, 1) \cdot k = [0, k)$ , and  $i$  is the set of indices between  $[0, k]$ . The sequence (13) will be negative until  $i > \lfloor uk \rfloor$ , and since  $\lfloor uk \rfloor < uk$ , we are ensured there is only one sign change.

Since the coefficients of (8) undergo one sign change, we again invoke Descartes' rule, and observe that we have one real positive root.  $\square$

## .3 Proof of Theorem 1

*Proof.* Let  $x = ku$ . Then we can think of (8) as

$$F(y, x) = \left(\frac{x}{k!} - \frac{1}{(k-1)!}\right)y^k + \dots + \left(\frac{x}{2!} - 1\right)y^2 + (x - 1)y + x \quad (14)$$

Implicit differentiation of (14), written as  $D_x F + D_y F \cdot y'$  where  $y' = dy/dx$ , gives

$$0 = \left(\frac{y^k}{k!} + \dots + y + 1\right) + \left(\left(\frac{1}{(k-1)!} - \frac{x}{k!}\right)ky^{k-1} + \dots + (1 - x)\right)y' \quad (15)$$

Noting that  $(D_x F)(y) = \frac{y^k}{k!} + \dots + y + 1$  and  $(D_y F)(x, y) = \left(\frac{1}{(k-1)!} - \frac{x}{k!}\right)ky^{k-1} + \dots + (1 - x)$  so that

$$y' = -D_x F \cdot (D_y F)^{-1} \quad (16)$$

**Proposition 1.** *Let  $(x, y)$  be a positive solution to  $F(x, y) = 0$ , then  $y'$  evaluated at that solution is positive.*

We first show the theorem assuming the proposition is true. We can similarly compute the second order implicit derivative  $d^2y/dx^2$ ; indeed,

$$y'' = \frac{D_x F \cdot (D_y^2 F \cdot y' + D_{x,y} F) - D_y F \cdot D_{y,x} F \cdot y'}{(D_y F)^2} \quad (17)$$

Hence, if  $D_x F \cdot (D_y^2 F \cdot y' + D_{x,y} F) - D_y F \cdot D_{y,x} F \cdot y' > 0$  then  $y'' > 0$ . We have

$$D_x F \cdot (D_y^2 F \cdot (-D_x F \cdot (D_y F)^{-1}) + \quad (18)$$

$$D_{x,y} F) - D_y F \cdot D_{y,x} F \cdot (-D_x F \cdot (D_y F)^{-1}) \\ = D_x F \cdot (D_y^2 F \cdot (-D_x F \cdot (D_y F)^{-1}) + 2D_{y,x} F) \quad (19)$$

$$= D_x F \cdot h(x, y) \quad (20)$$

where  $h(x, y) = D_y^2 F \cdot y' + 2D_{y,x} F$ . Since  $D_x F > 0$ , we focus on  $h(x, y)$ : Now,

$$(D_{y,x} F)(y) = ((k-1)!)^{-1} y^{k-1} + \dots + 1 \quad (21)$$

and

$$-D_y^2 F = \left(\frac{x}{k!} - \frac{1}{(k-1)!}\right) k(k-1) y^{k-2} + \dots + 2\left(\frac{x}{2} - 1\right) \quad (22)$$

Collecting all the  $x$  terms in  $D_y^2 F$  we can define

$$\tilde{h}(x, y) = \frac{x}{(k-2)!} y^{k-2} + \dots + x. \quad (23)$$

Since  $F(y, x) = 0$ , we have

$$\frac{x}{k!} y^k + \frac{x}{(k-1)!} y^{k-1} + \dots + x = \frac{1}{(k-1)!} y^k + \dots + y \quad (24)$$

so that

$$\tilde{h}(x, y) + \frac{x}{k!} y^k + \frac{x}{(k-1)!} y^{k-1} - \frac{x}{k!} y^k - \frac{x}{(k-1)!} y^{k-1} \\ = \frac{1}{(k-1)!} y^k + \dots + y - \frac{x}{k!} y^k - \frac{x}{(k-1)!} y^{k-1}$$

Then,

$$D_y^2 F = \frac{x}{k!} y^k + \frac{x}{(k-1)!} y^{k-1} + \frac{k}{(k-2)!} y^{k-2} + \dots + 2 \\ - \frac{1}{(k-1)!} y^k - \dots - y.$$

so that

$$h(x, y) = \frac{2}{(k-1)!} y^{k-1} + \dots + 2 - \left( \frac{1}{(k-1)!} y^k + \dots + y - \frac{x}{k!} y^k \right. \\ \left. - \frac{x}{(k-1)!} y^{k-1} - \frac{k}{(k-2)!} y^{k-2} - \dots - 2 \right) y' \\ = y' \left( \frac{x}{k!} - \frac{1}{(k-1)!} \right) y^k \\ + y' \left( \frac{2}{(k-1)! y'} + \frac{x}{(k-1)!} - \frac{1}{(k-2)!} \right) y^{k-1} \\ + y' \left( \frac{2}{(k-2)! y'} + \frac{k}{(k-2)!} - \frac{1}{(k-3)!} \right) y^{k-2} \\ + y' \left( \frac{2}{(k-3)! y'} + \frac{k-1}{(k-3)!} - \frac{1}{(k-4)!} \right) y^{k-3} \\ \vdots \\ + y' \left( \frac{2}{y'} + 2 \right).$$



Through straightforward, but somewhat cumbersome algebra, we can show that if  $(x, y)$  is a pair such that  $F(x, y) = 0$ , then

$$\frac{2}{y'} + 1 \geq x.$$

Following the above inequalities and using  $\frac{2}{y'} + 2 \geq x$ , at the solution  $(x, y)$  where  $F(x, y) = 0$

$$\begin{aligned} h(x, y) &\geq y' \left( \frac{x}{k!} - \frac{1}{(k-1)!} \right) y^k \\ &\quad + y' \left( \frac{x}{(k-1)!} - \frac{1}{(k-2)!} \right) y^{k-1} \\ &\quad + y' \left( \frac{x}{(k-2)!} - \frac{1}{(k-3)!} \right) y^{k-2} \\ &\quad \vdots \\ &\quad + y'(x) \\ &= y'F(x, y) \\ &= 0, \end{aligned}$$

and  $y'' \geq 0$  follows from  $h(x, y) \geq 0$ .

Now we prove Prop. 1. This lemma follows from the Gauss-Lucas Theorem [Meserve, 1982b], which states that if  $p(z)$  is a polynomial with real coefficients with complex roots  $r_1, \dots, r_n$ , then the complex roots of  $p'(z)$  is contained in the convex hull of  $r_1, \dots, r_n$ . For a fix  $x$ , applying this theorem to  $D_y F$  yields the fact that real parts of all roots of  $D_y F$  is less than the root of  $F(x, y)$ . Since  $D_y F \rightarrow -\infty$  as  $y \rightarrow \infty$ , at the root of  $F(x, y)$ ,  $D_y F \leq 0$ . By (16) and the fact  $D_x F > 0$ ,  $y' > 0$ .  $\square$